


WHAT IS CLAIMED IS:

1. A compound of structural formula I:



a pharmaceutically acceptable salt or a stereoisomer thereof,

wherein:

a and b are each independently chosen from a double bond and a single bond;

X is hydrogen or halogen;

10 when a is a single bond, Y and Z are each independently chosen from hydrogen, C₁₋₄ alkyl, and halogen,

or Y and Z, together with the carbon atom to which they are attached, form a cyclopropyl group;

when a is a double bond, Y is chosen from hydrogen, C₁₋₄ alkyl, and halogen;

n is 0, 1, 2, or 3;

U, V, W, and D are each independently chosen from CH, N, S, and O, provided that at least one of U, V,

15 W, and D is chosen from N, S, and O, and further provided that when one of U, V, W,

and D is S or O, then the other ring members are independently chosen from N and CH;

R¹ is chosen from hydrogen, CF₃, carbonyl(C₁₋₃ alkyl), hydroxyl, C₁₋₄ alkoxy, halogen, C₁₋₃ alkyl,

hydroxymethyl, and (C₀₋₆ alkyl)₂amino, wherein said alkyl and alkoxy are each optionally

substituted with one to seven fluorine atoms;

20 R² is chosen from:

halogen,

(carbonyl)₀₋₁C₁₋₁₀ alkyl,

(carbonyl)₀₋₁C₂₋₁₀ alkenyl,

(carbonyl)₀₋₁C₂₋₁₀ alkynyl,

25 C₁₋₁₀ alkenylamino,

- (carbonyl)0-1aryl C0-10 alkyl,
C3-8 cycloalkyl C0-10 alkyl,
(C3-8)heterocyclyl C0-10 alkyl,
C3-8 heterocycloalkyl C0-10 alkyl,
5 C1-4acylamino C0-10 alkyl,
C0-10 alkylamino C0-10 alkyl,
di-(C1-10 alkyl)amino C0-10 alkyl,
arylC0-10 alkylamino C0-10 alkyl,
(arylC0-10 alkyl)2amino C0-10 alkyl,
10 C3-8 cycloalkyl C0-10 alkylamino C0-10 alkyl,
C3-8 heterocyclyl C0-10 alkylamino C0-10 alkyl,
C3-8 heterocycloalkyl C0-10 alkylamino C0-10 alkyl,
(C3-8 cycloalkyl C0-10 alkyl)2amino C0-10 alkyl,
(C3-8 heterocyclyl C0-10 alkyl)2amino C0-10 alkyl,
15 (C3-8 heterocycloalkyl C0-10 alkyl)2amino C0-10 alkyl,
C3-8 cycloalkyl C0-10 alkyl aminocarbonylamino,
(C1-10 alkyl)2aminocarbonylamino,
(aryl C1-10 alkyl)1-2aminocarbonylamino,
C0-10 alkyl aminocarbonylamino,
20 C3-8 heterocyclyl C0-10 alkyl aminocarbonylamino,
C3-8 heterocycloalkyl C0-10 alkyl aminocarbonylamino,
(C1-10 alkyl)2aminocarbonyl C0-10 alkyl,
(aryl C1-10 alkyl)1-2aminocarbonyl C0-10 alkyl,
C0-10 alkyl aminocarbonyl C0-10 alkyl,
25 C3-8 cycloalkyl C0-10 alkyl aminocarbonyl C0-10 alkyl,
C3-8 heterocyclyl C0-10 alkyl aminocarbonyl C0-10 alkyl,
C3-8 heterocycloalkyl C0-10 alkyl aminocarbonyl C0-10 alkyl,
aryl C0-10 alkyl aminocarbonyl C0-10 alkyl,
C0-10 alkyl carbonylamino C0-10 alkyl,
30 C3-8 cycloalkyl C0-10 alkyl carbonylamino C0-10 alkyl,
C3-8 heterocyclyl C0-10 alkyl carbonylamino C0-10 alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 amino C₀₋₁₀ alkyl carbimidoyl C₀₋₁₀ alkylamino,
 (C₁₋₁₀ alkyl)₂aminocarbonyl,
 5 (aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonyl,
 C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
 C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
 carboxy C₀₋₁₀ alkyl,
 carboxy aryl,
 10 carboxy C₃₋₈ cycloalkyl,
 carboxy C₃₋₈ heterocyclyl,
 carboxy C₃₋₈ heterocycloalkyl,
 C₁₋₁₀ alkoxy,
 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 15 aryloxy,
 C₃₋₈ cycloalkyloxy,
 C₃₋₈ heterocyclyloxy,
 C₃₋₈ heterocycloalkyloxy,
 C₁₋₁₀ alkylcarbonyloxy,
 20 C₃₋₈ heterocyclyl C₀₋₁₀ alkylcarbonyloxy,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 aryl C₀₋₁₀ alkylcarbonyloxy,
 C₁₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 25 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 30 (aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,

- (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ cycloalkyl C₀₋₁₀alkyl)₁₋₂aminocarbonyloxy,
 hydroxy C₀₋₁₀alkyl,
 hydroxycarbonylC₀₋₁₀alkoxy,
 5 hydroxycarbonylC₀₋₁₀alkyloxy,
 C₁₋₁₀ alkylthio,
 C₁₋₁₀ alkylsulfinyl,
 aryl C₀₋₁₀ alkylsulfinyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfinyl,
 10 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfinyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfinyl,
 C₁₋₁₀ alkylsulfonyl,
 aryl C₀₋₁₀ alkylsulfonyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfonyl,
 15 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfonyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfonyl,
 C₁₋₁₀ alkylsulfonylamino,
 aryl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
 20 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
 cyano,
 nitro,
 perfluoroC₁₋₆alkyl, and
 25 perfluoroC₁₋₆alkoxy, and


wherein R² is optionally substituted with at least one substituent, R³, chosen from:

- halogen,
 (carbonyl)₀₋₁C₁₋₁₀ alkyl,
 (carbonyl)₀₋₁C₂₋₁₀ alkenyl,
 30 (carbonyl)₀₋₁C₂₋₁₀ alkynyl,
 (carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocycloalkyl C₀₋₁₀ alkyl,
C₁₋₄acylamino C₀₋₁₀ alkyl,
5 C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,
arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
(arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
10 C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
C₀₋₁₀ alkyl carbimidoylC₀₋₁₀ alkyl,
(C₁₋₁₀ alkyl)₂aminocarbonyl,
C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
15 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
(C₁₋₁₀ alkyl)₂aminocarbonyloxy,
hydroxycarbonylC₀₋₁₀alkoxy,
(C₁₋₁₀ alkyl)₂aminocarbonyloxy,
(aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
20 hydroxy C₀₋₁₀alkyl,
C₁₋₁₀ alkylsulfonyl,
C₁₋₁₀ alkylsulfonylamino,
aryl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
25 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
cyano,
nitro,
perfluoroC₁₋₆alkyl, and
30 perfluoroC₁₋₆alkoxy,

wherein R³ is optionally substituted with one or more groups chosen from hydrogen, OH, (C₁₋₆)alkoxy, halogen, CO₂H, CN, O(C=O)C₁₋₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl, and NH₂.

2. A compound according to Claim 1, wherein X is fluorine.
3. A compound according to Claim 1, wherein X is hydrogen.
4. A compound according to Claim 1, wherein a is a single bond and b is a double
5. A compound according to Claim 1 and of structural formula II, wherein:



a pharmaceutically acceptable salt or a stereoisomer thereof,

wherein:


a and b are each independently chosen from a double bond and a single bond;

n is 0, 1, 2, or 3;


X is hydrogen or halogen;

when a is a single bond, Y and Z are each independently chosen from hydrogen, C₁₋₄ alkyl, and halogen,
or Y and Z, together with the carbon atom to which they are attached, form a cyclopropyl
group;

when a is a double bond, Y is chosen from hydrogen, C₁₋₄ alkyl, and halogen;



is chosen from:



E is S or O;

R¹ is chosen from: hydrogen, CF₃, carbonyl(C₁₋₃ alkyl), hydroxyl, C₁₋₄ alkoxy, halogen, C₁₋₃ alkyl, hydroxymethyl, and (C₀₋₆ alkyl)₂amino, wherein said alkyl and alkoxy are each optionally substituted with one to seven fluorine atoms;

R² is chosen from:

halogen,

(carbonyl)₀₋₁C₁₋₁₀ alkyl,

(carbonyl)₀₋₁C₂₋₁₀ alkenyl,

(carbonyl)₀₋₁C₂₋₁₀ alkynyl,

C₁₋₁₀ alkenylamino,

(carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,

(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl,

C₁₋₄acylamino C₀₋₁₀ alkyl,

C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,

arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

(arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,

C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,
 5 (C₁₋₁₀ alkyl)₂aminocarbonylamino,
 (aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonylamino,
 C₀₋₁₀ alkyl aminocarbonylamino,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,
 10 (C₁₋₁₀ alkyl)₂aminocarbonyl C₀₋₁₀ alkyl,
 (aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonyl C₀₋₁₀ alkyl,
 C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
 15 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
 aryl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
 C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 20 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 amino C₀₋₁₀ alkyl carbimidoylC₀₋₁₀ alkylamino,
 (C₁₋₁₀ alkyl)₂aminocarbonyl,
 (aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonyl,
 25 C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
 C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
 carboxy C₀₋₁₀ alkyl,
 carboxy aryl,
 carboxy C₃₋₈ cycloalkyl,
 30 carboxy C₃₋₈ heterocyclyl,
 carboxy C₃₋₈ heterocycloalkyl,

- C₁₋₁₀ alkoxy,
 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 aryloxy,
 C₃₋₈ cycloalkyloxy,
 5 C₃₋₈ heterocyclyloxy,
 C₃₋₈ heterocycloalkyloxy,
 C₁₋₁₀ alkylcarbonyloxy,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkylcarbonyloxy,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 10 C₃₋₈ cycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 aryl C₀₋₁₀ alkylcarbonyloxy,
 C₁₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 15 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 (aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 20 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ cycloalkyl C₀₋₁₀alkyl)₁₋₂aminocarbonyloxy,
 hydroxy C₀₋₁₀alkyl,
 hydroxycarbonylC₀₋₁₀alkoxy,
 hydroxycarbonylC₀₋₁₀alkyloxy,
 25 C₁₋₁₀ alkylthio,
 C₁₋₁₀ alkylsulfinyl,
 aryl C₀₋₁₀ alkylsulfinyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfinyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfinyl,
 30 C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfinyl,
 C₁₋₁₀ alkylsulfonyl,

aryl C₀₋₁₀ alkylsulfonyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfonyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfonyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfonyl,
5 C₁₋₁₀ alkylsulfonylamino,
aryl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
10 cyano,
nitro,
perfluoroC₁₋₆alkyl, and
perfluoroC₁₋₆alkoxy, and

wherein R² is optionally substituted with at least one substituent R³;

15 R³ is chosen from:

halogen,
(carbonyl)₀₋₁C₁₋₁₀ alkyl,
(carbonyl)₀₋₁C₂₋₁₀ alkenyl,
(carbonyl)₀₋₁C₂₋₁₀ alkynyl,
20 (carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocycloalkyl C₀₋₁₀ alkyl,
C₁₋₄acylamino C₀₋₁₀ alkyl,
25 C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,
arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
(arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
30 C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

- C₀₋₁₀ alkyl carbimidoylC₀₋₁₀ alkyl,
 (C₁₋₁₀ alkyl)₂aminocarbonyl,
 C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 5 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 hydroxycarbonylC₀₋₁₀alkoxy,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 (aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 hydroxy C₀₋₁₀alkyl,
 10 C₁₋₁₀ alkylsulfonyl,
 C₁₋₁₀ alkylsulfonylamino,
 aryl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
 15 C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
 cyano,
 nitro,
 perfluoroC₁₋₆alkyl, and
 perfluoroC₁₋₆alkoxy,
 20 wherein R³ is optionally substituted with one or more groups chosen from hydrogen, OH, (C₁₋₆)alkoxy,
 halogen, CO₂H, CN, O(C=O)C₁₋₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl, and NH₂.

6. A compound according to Claim 5, wherein R¹ is chosen from: hydrogen, CF₃,
 25 hydroxyl, and C₁₋₃ alkyl optionally substituted with one to seven fluorine atoms.

7. A compound according to Claim 6, wherein R¹ is chosen from: hydrogen and
 C₁₋₃ alkyl.

8. A compound according to Claim 7, wherein R¹ is methyl.

9. A compound according to Claim 8, wherein R² is chosen from:

halogen,

(carbonyl)₀₋₁C₁₋₁₀ alkyl,

(carbonyl)₀₋₁C₂₋₁₀ alkenyl,

5 (carbonyl)₀₋₁C₂₋₁₀ alkynyl,

C₁₋₁₀ alkenylamino,

(carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,

(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,

10 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl,

C₁₋₄ acylamino C₀₋₁₀ alkyl,

C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,

arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

15 (arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,

20 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,

(C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,

(C₁₋₁₀ alkyl)₂aminocarbonylamino,

(aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonylamino,

25 C₀₋₁₀ alkyl aminocarbonylamino,

C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonylamino,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,

C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,


30 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,



C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
 amino C₀₋₁₀ alkyl carbimidoyl C₀₋₁₀ alkylamino,
 C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
 C₁₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 5 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 C₁₋₁₀ alkylsulfonylamino,
 10 aryl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
 cyano,
 15 nitro,
 perfluoroC₁₋₆alkyl, and
 perfluoroC₁₋₆alkoxy, and


wherein R² is optionally substituted with at least one substituent R³.

10. A compound according to Claim 9, wherein E is S.

11. A compound according to Claim 10, wherein  is:




and



12. A compound according to Claim 11, wherein b is a double bond.

13. A compound according to Claim 12, wherein a is a single bond and b is a double bond.

14. A compound according to Claim 1 and of structural formula III,



a pharmaceutically acceptable salt or a stereoisomer thereof, wherein:

X is hydrogen or halogen;

10 n is 0, 1, 2, or 3;

Y and Z are each independently chosen from hydrogen, C₁₋₄ alkyl, and halogen, or Y and Z, together with the carbon atom to which they are attached, form a cyclopropyl group;

U, V, W, and D are each independently chosen from N and CH, provided that at least one of U, V, W, and D is CH;

15 R² is chosen from:

halogen,

(carbonyl)₀₋₁C₁₋₁₀ alkyl,

(carbonyl)₀₋₁C₂₋₁₀ alkenyl,

(carbonyl)₀₋₁C₂₋₁₀ alkynyl,

20 C₁₋₁₀ alkenylamino,

(carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,

(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl,

25 C₁₋₄acylamino C₀₋₁₀ alkyl,

- C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,
arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
(arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
5 C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
(C₃₋₈ cycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
(C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
10 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,
(C₁₋₁₀ alkyl)₂aminocarbonylamino,
(aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonylamino,
C₀₋₁₀ alkyl aminocarbonylamino,
15 C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonylamino,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,
(C₁₋₁₀ alkyl)₂aminocarbonyl C₀₋₁₀ alkyl,
(aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonyl C₀₋₁₀ alkyl,
C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
20 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
aryl C₀₋₁₀ alkyl aminocarbonyl C₀₋₁₀ alkyl,
C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
25 C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,
amino C₀₋₁₀ alkyl carbimidoylC₀₋₁₀ alkylamino,
30 (C₁₋₁₀ alkyl)₂aminocarbonyl,
(aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonyl,

C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
 C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,
 carboxy C₀₋₁₀ alkyl,
 carboxy aryl,
 5 carboxy C₃₋₈ cycloalkyl,
 carboxy C₃₋₈ heterocyclyl,
 carboxy C₃₋₈ heterocycloalkyl,
 C₁₋₁₀ alkoxy,
 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 10 aryloxy,
 C₃₋₈ cycloalkyloxy,
 C₃₋₈ heterocycliloxy,
 C₃₋₈ heterocycloalkyloxy,
 C₁₋₁₀ alkylcarbonyloxy,
 15 C₃₋₈ heterocyclyl C₀₋₁₀ alkylcarbonyloxy,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylcarbonyloxy,
 aryl C₀₋₁₀ alkylcarbonyloxy,
 C₁₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 20 C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 25 (aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ heterocyclyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 (C₃₋₈ cycloalkyl C₀₋₁₀alkyl)₁₋₂aminocarbonyloxy,
 hydroxy C₀₋₁₀alkyl,
 30 hydroxycarbonylC₀₋₁₀alkoxy,
 hydroxycarbonylC₀₋₁₀alkyloxy,

C₁₋₁₀ alkylthio,
C₁₋₁₀ alkylsulfinyl,
aryl C₀₋₁₀ alkylsulfinyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfinyl,
5 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfinyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfinyl,
C₁₋₁₀ alkylsulfonyl,
aryl C₀₋₁₀ alkylsulfonyl,
C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfonyl,
10 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfonyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfonyl,
C₁₋₁₀ alkylsulfonylamino,
aryl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
15 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
cyano,
nitro,
perfluoroC₁₋₆alkyl, and
20 perfluoroC₁₋₆alkoxy, and

wherein R² is optionally substituted with at least one substituent, R³, chosen from:

halogen,
(carbonyl)₀₋₁C₁₋₁₀ alkyl,
(carbonyl)₀₋₁C₂₋₁₀ alkenyl,
25 (carbonyl)₀₋₁C₂₋₁₀ alkynyl,
(carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,
C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,
(C₃₋₈)heterocycloalkyl C₀₋₁₀ alkyl,
30 C₁₋₄acylamino C₀₋₁₀ alkyl,
C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

di-(C₁₋₁₀ alkyl)amino C₀₋₁₀ alkyl,
 arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
 (arylC₀₋₁₀ alkyl)₂amino C₀₋₁₀ alkyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
 5 C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,
 C₀₋₁₀ alkyl carbimidoylC₀₋₁₀ alkyl,
 (C₁₋₁₀ alkyl)₂aminocarbonyl,
 C₁₋₁₀ alkoxy (carbonyl)₀₋₁C₀₋₁₀ alkyl,
 10 C₁₋₁₀alkyloxy C₀₋₁₀alkyl,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 hydroxycarbonylC₀₋₁₀alkoxy,
 (C₁₋₁₀ alkyl)₂aminocarbonyloxy,
 (aryl C₀₋₁₀ alkyl)₁₋₂aminocarbonyloxy,
 15 hydroxy C₀₋₁₀alkyl,
 C₁₋₁₀ alkylsulfonyl,
 C₁₋₁₀ alkylsulfonylamino,
 aryl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
 20 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
 cyano,
 nitro,
 perfluoroC₁₋₆alkyl, and
 25 perfluoroC₁₋₆alkoxy, and

wherein R³ is optionally substituted with one or more groups chosen from hydrogen, OH, (C₁₋₆)alkoxy,
 halogen, CO₂H, CN, O(C=O)C₁₋₆ alkyl, NO₂, trifluoromethoxy, trifluoroethoxy, -O(0-1)(C₁₋₁₀)perfluoroalkyl, and NH₂.

30 15. A compound according to Claim 14, wherein X is hydrogen.

16. A compound according to Claim 15, wherein R² is chosen from:

halogen,

(carbonyl)₀₋₁C₁₋₁₀ alkyl,

(carbonyl)₀₋₁C₂₋₁₀ alkenyl,

(carbonyl)₀₋₁aryl C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl,

(C₃₋₈)heterocyclyl C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl,

C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

arylC₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocyclyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylamino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,

(aryl C₁₋₁₀ alkyl)₁₋₂aminocarbonylamino,

C₀₋₁₀ alkyl aminocarbonylamino,

C₃₋₈ heterocyclyl C₀₋₁₀ alkyl aminocarbonylamino,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl aminocarbonylamino,

C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

C₃₋₈ cycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocyclyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

aryl C₀₋₁₀ alkyl carbonylamino C₀₋₁₀ alkyl,

C₀₋₁₀ alkylcarboxy C₀₋₁₀ alkylamino,

C₁₋₁₀ alkoxy,

C₁₋₁₀alkyloxy C₀₋₁₀alkyl,

aryloxy,

C₃₋₈ cycloalkyloxy,

C₃₋₈ heterocyclioxy,

C₃₋₈ heterocycloalkyloxy,

C₁₋₁₀ alkyloxy(carbonyl)₀₋₁C₀₋₁₀ alkylamino,

C₃₋₈ heterocyclyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 aryl C₀₋₁₀ alkyloxy(carbonyl)₀₋₁ C₀₋₁₀ alkylamino,
 5 hydroxy C₀₋₁₀alkyl,
 C₁₋₁₀ alkylthio,
 C₁₋₁₀ alkylsulfonyl,
 aryl C₀₋₁₀ alkylsulfonyl,
 C₃₋₈ heterocyclyl C₀₋₁₀ alkylsulfonyl,
 10 C₃₋₈ heterocycloalkyl C₀₋₁₀ alkylsulfonyl,
 C₃₋₈ cycloalkyl C₀₋₁₀ alkylsulfonyl,
 C₁₋₁₀ alkylsulfonylamino,
 aryl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ heterocyclyl C₁₋₁₀ alkylsulfonylamino,
 15 C₃₋₈ heterocycloalkyl C₁₋₁₀ alkylsulfonylamino,
 C₃₋₈ cycloalkyl C₁₋₁₀ alkylsulfonylamino,
 cyano,
 nitro,
 perfluoroC₁₋₆alkyl, and
 20 perfluoroC₁₋₆alkoxy, and

wherein R² is optionally substituted with at least one substituent R³.

17. A compound according to Claim 16, wherein at least two of U, V, W, and D are each N and provided that at least one of U, V, W, and D is CH

25

18. A compound according to Claim 1, selected from:

17β-[2-(butylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(anilino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(pyridin-2-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 30 17β-[(2-methylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[[2-methyl(phenyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[[2-(4-fluorophenyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;

- 17 β -[2-(benzylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(isopropylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyridin-3-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(2-fluorophenyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
5 17 β -[2-(methoxyethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(2-piperid-1-ylethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(t-butyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(4-cyanophenyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(cyclohexyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
10 17 β -[2-[(pyridin-4-ylmethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyrimidin-2-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyridin-4-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(cyclopropylmethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(propylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
15 17 β -[2-(allylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(heptylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(octylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(hexylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(5-methyl-1,2,3-thiadiazol-2-yl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
20 17 β -[2-(methoxypropyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(2-morpholin-1-ylethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(2,2,2-trifluoroethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-[(pyridin-2-ylethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(amino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
25 17 β -[2-(guanidino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(1-methyl-1H-imidazole-5-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(acetamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(phenyl carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(thiophene-3-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
30 17 β -[2-(furan-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyrazine-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyridine-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(thiophene-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
17 β -[2-(pyridine-3-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;

- 17β-[2-(pyridine-4-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(1-*t*-butyl-3-methyl-1H-pyrazole-5-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(1-methyl-proline-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 5 17β-[2-(1-methyl-1H-imidazole-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(1H-imidazole-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(methanesulfonamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(ethyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(isopropyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 10 17β-[2-(2-fluoroethyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(*t*-butylcarbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one
 17β-[2-(ureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(N'-pyridin-2-ylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(N'-cyclopropylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 15 17β-[2-(N'-cyclohexylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(N'-cyclohexylmethylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(morpholine-4-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(piperazine-1-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[N'-isopropylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 20 17β-[2-(pyridyl-3-ureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-[N'-(methylamino)ethylureyl]-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(ureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androstan-3-one;
 17β-[2-(pyridin-2-yl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androstan-3-one;
 17β-[2-(methyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 25 17β-[2-(pyrid-3-yl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(ethyl acetyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(acetonitril-yl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(2-chlorophenyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(methyl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 30 17β-[2-(phenyl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(3,5-dimethylpyrazol-1-yl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[2-(aminoacetyl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[5-(amino)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;
 17β-[5-(ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5α-androst-1-en-3-one;

17 β -[5-(N-methyl-ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[5-(N,N-dimethyl-ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one; and
 pharmaceutically acceptable salts and stereoisomers thereof.

- 5 19. A compound according to Claim 18, selected from:
- 17 β -[2-(butylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(anilino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridin-2-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(methylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 10 17 β -{[2-methyl(phenyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{[2-(4-fluorophenyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(benzylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(isopropylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridin-3-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 15 17 β -{[2-(2-fluorophenyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{[2-(methoxyethyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-[(2-piperid-1-ylethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{[2-(t-butyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{[2-(4-cyanophenyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 20 17 β -[2-[(cyclohexyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-[(pyridin-4-ylmethyl)amino]-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyrimidin-2-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridin-4-ylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{2-[(cyclopropylmethyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 25 17 β -[2-(propylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(allylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(heptylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(octylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(hexylamino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 30 17 β -{2-[(5-methyl-1,2,3-thiadiazol-2-yl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{[2-(methoxypropyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{2-[(2-morpholin-1-ylethyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{2-[(2,2,2-trifluoroethyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -{2-[(pyridin-2-ylethyl)amino]-1,3-thiazol-4-yl}-4-methyl-4-aza-5 α -androst-1-en-3-one;

- 17 β -[2-(amino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(guanidino)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(1-methyl-1H-imidazole-5-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(acetamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 5 17 β -[2-(phenyl carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(thiophene-3-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(furan-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyrazine-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridine-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 10 17 β -[2-(thiophene-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridine-3-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridine-4-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(1-*t*-butyl-3-methyl-1H-pyrazole-5-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 15 17 β -[2-(1-methyl-proline-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(1-methyl-1H-imidazole-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(1H-imidazole-2-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(methanesulfonamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(ethyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 20 17 β -[2-(isopropyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(2-fluoroethyl carbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(*t*-butylcarbamate)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(ureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(N'-pyridin-2-ylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 25 17 β -[2-(N'-cyclopropylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(N'-cyclohexylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(N'-cyclohexylmethylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(morpholine-4-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(piperazine-1-carboxamido)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 30 17 β -[N'-isopropylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridin-3-ylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-[N'-(methylamino)ethylureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(ureyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyridin-2-yl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;

17 β -[2-(methyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(pyrid-3-yl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(ethyl acetyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(acetonitrilyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 5 17 β -[2-(2-chlorophenyl)-1,3-thiazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one; and
 pharmaceutically acceptable salts and stereoisomers thereof.

20. A compound according to Claim 19, selected from: 17 β -[2-(methyl)-1,3-
 imidazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 10 17 β -[2-(phenyl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(3,5-dimethylpyrazol-1-yl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[2-(aminoacetyl)-1,3-imidazol-4-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[5-(amino)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[5-(ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 15 17 β -[5-(N-methyl-ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one;
 17 β -[5-(N,N-dimethyl-ureyl)-1,2,4-triazol-3-yl]-4-methyl-4-aza-5 α -androst-1-en-3-one; and
 pharmaceutically acceptable salts and stereoisomers thereof.

21. A method for modulating a function mediated by the androgen receptor in a
 20 mammal in need of such modulation comprising administering a therapeutically effective amount of a
 compound of Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

22. A method of activating the function of the androgen receptor in a mammal in
 need of such activation comprising administering a therapeutically effective amount of a compound of
 25 Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

23. A method of Claim 22, wherein said function mediated by the androgen receptor
 is activated in bone or muscle tissue and blocked in the prostate or the uterus.

24. A method of treating a condition in a mammal which is caused by androgen
 30 deficiency, which can be ameliorated by androgen replacement, or which can be increased by androgen
 replacement, which condition is selected from weakened muscle tone, osteoporosis, osteopenia,
 glucocorticoid-induced osteoporosis, periodontal disease, bone fracture, bone damage following bone
 reconstructive surgery, sarcopenia, frailty, aging skin, male hypogonadism, postmenopausal symptoms in

women, atherosclerosis, hypercholesterolemia, hyperlipidemia, obesity, aplastic anemia and other hematopoietic disorders, inflammatory arthritis and joint repair, HIV-wasting, prostate cancer, benign prostatic hyperplasia (BPH), abdominal adiposity, metabolic syndrome, type II diabetes, cancer cachexia, Alzheimer's disease, muscular dystrophies, cognitive decline, sexual dysfunction, sleep apnea, depression, premature ovarian failure, and autoimmune disease, comprising administering to the mammal in need of such treatment, a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

25. A method according to Claim 24, wherein said condition is osteoporosis.

26. A method of treating osteoporosis in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

27. A method of Claim 26, further comprising the administration of an agent selected from:

- 1) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative,
- 2) a bisphosphonate,
- 3) an antiestrogen or a selective estrogen receptor modulator,
- 4) an $\alpha v \beta 3$ integrin receptor antagonist,
- 5) a cathepsin K inhibitor,
- 6) an HMG-CoA reductase inhibitor,
- 7) an osteoclast vacuolar ATPase inhibitor,
- 8) an antagonist of VEGF binding to osteoclast receptors,
- 9) an activator of peroxisome proliferator-activated receptor γ ,
- 10) calcitonin,
- 11) a calcium receptor antagonist,
- 12) parathyroid hormone or analog thereof,
- 13) a growth hormone secretagogue,
- 14) human growth hormone,
- 15) insulin-like growth factor,
- 16) a p38 protein kinase inhibitor,

- 17) bone morphogenetic protein,
- 18) an inhibitor of BMP antagonism,
- 19) a prostaglandin derivative,
- 20) vitamin D or vitamin D derivative,
- 21) vitamin K or vitamin K derivative,
- 22) ipriflavone,
- 23) fluoride salts,
- 24) dietary calcium supplement, and
- 25) osteoprotegerin.

28. The method according to Claim 27, wherein:

- 1) the estrogen or estrogen derivative, alone or in combination with a progestin or progestin derivative, is selected from conjugated estrogen, equine estrogen, 17 β -estradiol, estrone, 17 β -ethynyl estradiol, 17 β -ethynyl estradiol with at least one agent selected from norethindrone and medroxyprogesterone acetate;
- 2) the bisphosphonate is selected from alendronate, clodronate, etidronate, ibandronate, incadronate, minodronate, neridronate, olpadronate, pamidronate, piridronate, risedronate, tiludronate, and zoledronate;
- 3) the antiestrogen or selective estrogen receptor modulator is selected from raloxifene, clomiphene, zuclomiphene, enclomiphene, nafoxidene, CI-680, CI-628, CN-55,945-27, Mer-25, U-11,555A, U-100A, tamoxifen, lasofoxifene, toremifene, azorxifene, EM-800, EM-652, TSE 424, droloxifene, idoxifene, and levormeloxifene;
- 4) the HMG-CoA reductase inhibitor is selected from lovastatin, simvastatin, dihydroxy-open acid simvastatin, pravastatin, fluvastatin, atorvastatin, cerivastatin, rosuvastatin, pitavastatin, and nisvastatin;
- 5) calcitonin is salmon calcitonin administered as a nasal spray;
- 6) bone morphogenetic protein is selected from BMP 2, BMP 3, BMP 5, BMP 6, BMP 7, TGF beta, and GDF5;
- 7) insulin-like growth factor is selected from IGF I and IGF II alone or in combination with IGF binding protein 3;
- 8) the prostaglandin derivative is selected from agonists of prostaglandin receptors EP₁, EP₂, EP₄, FP, and IP;
- 9) the fibroblast growth factor is selected from aFGF and bFGF;

10) parathyroid hormone (PTH) or PTH analog is selected from PTH subcutaneous injection, human PTH (1-84), human PTH (1-34), and other partial sequences, native or with substitutions;

11) vitamin D or vitamin D derivative is selected from natural vitamin D, 25-OH-vitamin D3, 1 α ,25(OH)₂ vitamin D3, 1 α -OH-vitamin D3, 1 α -OH-vitamin D2, dihydrotachysterol, 26,27-F6-1 α ,25(OH)₂ vitamin D3, 19-nor-1 α ,25(OH)₂ vitamin D3, 22-oxacalcitriol, calcipotriol, 1 α ,25(OH)₂-16-ene-23-yne-vitamin D3(Ro 23-7553), EB1089, 20-epi-1 α ,25(OH)₂ vitamin D3, KH1060, ED71, 1 α ,24(S)-(OH)₂ vitamin D3, and 1 α ,24(R)-(OH)₂ vitamin D3;

12) the dietary calcium supplement is selected from calcium carbonate, calcium citrate, and natural calcium salts; and

13) the fluoride salts are chosen from sodium fluoride and monosodium fluorophosphate (MFP); and pharmaceutically acceptable salts or stereoisomers thereof.

29. The method according to Claim 28, wherein the bisphosphonate is alendronate monosodium trihydrate or alendronate monosodium monohydrate.

30. The method of Claim 27, wherein said agent is selected from:

- 1) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative,
- 2) a bisphosphonate,
- 3) an antiestrogen or a selective estrogen receptor modulator,
- 4) an $\alpha v \beta 3$ integrin receptor antagonist,
- 5) a cathepsin K inhibitor,
- 6) an osteoclast vacuolar ATPase inhibitor,
- 7) calcitonin,
- 8) osteoprotegerin, and
- 9) parathyroid hormone or analog thereof.

31. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1 and a pharmaceutically acceptable carrier.

32. A composition of Claim 31, further comprising an active ingredient selected from:

- 1) an estrogen or an estrogen derivative, alone or in combination with a progestin or progestin derivative,
- 2) a bisphosphonate,
- 3) an antiestrogen or a selective estrogen receptor modulator,
- 4) an $\alpha v \beta 3$ integrin receptor antagonist,
- 5) a cathepsin K inhibitor,
- 6) an HMG-CoA reductase inhibitor,
- 7) an osteoclast vacuolar ATPase inhibitor,
- 8) an antagonist of VEGF binding to osteoclast receptors,
- 9) an activator of peroxisome proliferator-activated receptor γ ,
- 10) calcitonin,
- 11) a calcium receptor antagonist,
- 12) parathyroid hormone or analog thereof,
- 13) a growth hormone secretagogue,
- 14) human growth hormone,
- 15) insulin-like growth factor,
- 16) a p38 protein kinase inhibitor,
- 17) bone morphogenetic protein,
- 18) an inhibitor of BMP antagonism,
- 19) a prostaglandin derivative,
- 20) vitamin D or vitamin D derivative,
- 21) vitamin K or vitamin K derivative,
- 22) ipriflavone,
- 23) fluoride salts,
- 24) dietary calcium supplement, and
- 25) osteoprotegerin.

33. A composition of Claim 32, wherein said bisphosphonate is alendronate.

34. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

35. A method of increasing Bone Mineral Density in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

5

36. A method of reducing the risk of vertebral or non-vertebral fractures in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

10

37. A method of effecting a bone turnover marker in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof, wherein said bone turnover marker is selected from urinary C-telopeptide degradation products of type I collagen (CTX), urinary N-telopeptide cross-links of type I collagen (NTX), DXA, and DPD.

15

38. A pharmaceutical composition made by combining a compound according to Claim 1 and a pharmaceutically acceptable carrier.

20

39. A process for making a pharmaceutical composition comprising combining a compound according to Claim 1 and a pharmaceutically acceptable carrier.

25

40. A method of treating or preventing an arthritic condition in a mammal in need thereof, comprising administering a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt or a stereoisomer thereof.

41. A method of Claim 40, wherein the arthritic conditions is selected from rheumatoid arthritis and osteoarthritis.